# Pteropod molluscs Toward an understanding of population connectivity and species delimitations

Stephanie L. Bush<sup>1,2,3</sup>, Herman H. Wirshing<sup>3</sup>, Karen J. Osborn<sup>3</sup>, & Bruce H. Robison<sup>3</sup>

1 Monterey Bay Aquarium Research Institute, California, USA

2 Monterey Bay Aquarium, California, USA

3 Smithsonian Institution National Museum of Natural History, Washington DC, USA


#### DENSE POPULATIONS

#### Hyalocylis striata





#### CRITICAL FOOD WEB COMPONENTS



## ACIDIFICA



WARMING

Increase in metabolic rates



Increase demand for food

Increase in oxygen requirements


DEOXYGE

Metabolic suppression

May not be achievable in warmer water

#### Clio pyramidata

Hawaii





Southern California

Northwest Atlantic


(Bear Seamount)


#### Clio pyramidata






#### Cavolinia uncinata







#### Hyalocylis striata









#### Hyalocylis striata Fsтs



#### Corolla spectabilis





- Species delimitations depend on taxon
  - Behavioral, physiological, ecological, selection differences

- Cavolinia uncinata likely two species
  - Previously considered formae by van der Spoel

- Clio pyramidata cosmopolitan
  - Despite morphological differences and large geographical distances



### ACKNOWLED

Mike Vecchione -U.S. National Oceanic and Atmospheric Administration

Smithsonian Institute -National Museum of Natural History

Monterey Bay Aquarium Research Institute

U.S. UNOLS Chief Scientist Training Cruise